arXiv:cond-mat/0406350v1 [cond-mat.stat-mech] 16 Jun 2004

Breakdown of the perturbative renormalization group at certain quantum critical points

D. Belitz[1], T.R. Kirkpatrick[2], and J. Rollbühler[1]
[1]*Department of Physics and Materials Science Institute, University of Oregon, Eugene, OR 97403*
[2]*Institute for Physical Science and Technology, and Department of Physics*
University of Maryland, College Park, MD 20742

(Dated: February 2, 2008)

It is shown that the presence of multiple time scales at a quantum critical point can lead to a breakdown of the loop expansion for critical exponents, since coefficients in the expansion diverge. Consequently, results obtained from finite-order perturbative renormalization-group treatments may be not be an approximation in any sense to the true asymptotic critical behavior. This problem manifests itself as a non-renormalizable field theory, or, equivalently, as the presence of a dangerous irrelevant variable. The quantum ferromagnetic transition in disordered metals provides an example.

PACS numbers: 05.30.-d; 64.60.Ak; 64.60.Ht; 75.40.-s

The phenomenon known as critical slowing down is a dramatic effect near critical points [1]. It occurs because the order parameter (OP) relaxation rate τ diverges together with the correlation length ξ. The dynamical critical exponent z is defined by $\tau \propto \xi^z$. If soft modes other than the OP fluctuations couple to the latter, one needs to take them into account as well. For instance, at the liquid-gas critical point one needs to consider shear modes [1], and the quantum ferromagnetic transition [2, 3] is influenced by soft particle-hole excitations [4]. These additional soft modes are 'generic' in the sense that their softness is not related to the critical point (e.g., shear modes in a fluid are soft due to momentum conservation), and they introduce diverging time scales in addition to the critical one [5]. In contrast to the diverging length scale ξ, which is unique in any given direction, one therefore usually has to deal with multiple diverging time scales. In classical statistical mechanics this complication remains restricted to the dynamics, as the static critical behavior is independent of it, but in quantum statistical mechanics, where the statics and the dynamics are coupled, the critical behavior of thermodynamic quantities is also affected [2].

The technical description of phase transitions often relies on the renormalization group (RG). There are two different formulations of the RG. The first one, Wilson's momentum-shell RG [6, 7], integrates out degrees of freedom with successively larger wave numbers in order to derive a simpler effective theory valid only at asymptotically large length scales, i.e., at criticality. The second one, often referred to as the field-theoretic method, has its roots in high-energy physics. It introduces an artificial 'renormalized theory' in order to study the behavior of the physical theory under changes of the momentum cutoff, i.e., the basic length scale [8]. These two formulations of the RG are believed to be equivalent [9]. An important concept in the field-theoretic formulation is the 'renormalizability' of the theory, i.e., the possibility of constructing a renormalized theory that is finite in the limit of an infinite momentum cutoff by absorbing any infinities into a finite number of renormalization constants. Renormalizability is crucial in order to ensure scale independent RG flow equations and the power-law critical behavior that generically follows from them. While this concept has no obvious analog in the Wilsonian method, it is generally believed that lack of renormalizability can be remedied by adding additional renormalization constants, which in the Wilsonian method corresponds to the generation, under renormalization, of additional terms that were not present in the original action. In either method, one usually proceeds by performing a loop expansion in powers of a suitable coupling constant. In conjunction with an ϵ-expansion about some critical dimensionality, this allows for a perturbatively controlled determination of the critical behavior.

In this Letter we come to the remarkable conclusion that the perturbative RG can break down at a quantum critical point if there are multiple divergent time scales. Specifically, low-order results of an ϵ-expansion for critical exponents may be invalid because coefficients of terms that are formally of higher order in ϵ diverge. A second conclusion is that nonrenormalizability can be related to the existence of a dangerous irrelevant variable (DIV) [10] rather than to the generation of new terms in the momentum-shell method. Our results are generic, but for the sake of definiteness we illustrate them by using the quantum ferromagnetic transition in disordered metals [2] as an example. This transition is known to have an upper critical dimension $d_c^+ = 4$ [11, 12]. Accordingly, there is an effective quartic coupling constant u that is irrelevant with respect to a Gaussian mean-field fixed point (FP) in spatial dimensions $d > 4$, but becomes marginal in $d = 4$. Standard lore [6, 7] suggests that a controlled $\epsilon = 4 - d$ expansion of the critical exponents is possible for $d < 4$. Indeed, we will show below that to one-loop order one finds a flow equation

$$\frac{du}{d\ell} = \epsilon u - \frac{11}{12}\, u^2 + O(u^3), \tag{1}$$

where $\ell = \ln b$ with $b > 0$ the RG length scale factor. This suggests a FP value $u^* = 12\epsilon/11 + O(\epsilon^2)$, which determines the critical exponent η via $\eta = u^*/6 + O(\epsilon^2)$,

$$\eta = 2\epsilon/11 + O(\epsilon^2). \tag{2a}$$

The correlation length exponent ν and the dynamical exponent z are given in terms of η,

$$z = 4 - \eta \quad , \quad \nu = 1/(2-\eta). \tag{2b}$$

These results have nothing in common with the exact critial behavior found in Ref. 12, which is not even given by power laws. This creates an obvious conundrum, especially since the $4-\epsilon$ expansion is a more standard and straightforward method than the one employed in [12].

Equations (1,2), although formally correct, do not constitute an approximation to the true asymptotic critical behavior in any sense. This can first be seen in the flow equations at two-loop order, where the loop expansion parameter u couples to an irrelevant variable w,

$$\frac{du}{d\ell} = \epsilon u - \frac{11}{12}u^2 - \frac{u^3}{32}(1 - \ln w) + O(u^4), \tag{3a}$$

$$\frac{dw}{d\ell} = -2w + \frac{11}{12}uw + \frac{u^2}{32}(1 - \ln w)w + O(u^3). \tag{3b}$$

w is irrelevant with a scale dimension $-2 + O(\epsilon)$ and a FP value $w^* = 0$. It is common that some scaling function depends singularly on an irrelevant variable, making the latter a DIV with respect to the corresponding observable [10]. Here, however, the coefficients of the loop expansion itself depend singulary on w, leading to a breakdown of the perturbative RG [13]. This phenomenon is reminiscent of the breakdown of the virial expansion for transport coefficients in classical statistical mechanics [14]. In either case, a seemingly systematic expansion in powers of a coupling constant is ruined by the appearance of logarithmic terms. Physically, w reflects the ratio of the diffusive time scale of the itinerant electrons, which is characterized by a frequency scaling as a wave number squared, $\Omega \sim \boldsymbol{k}^2$, and the critical time scale characterized by $\Omega \sim \boldsymbol{k}^4$ [2, 15].

In the remainder of this Letter we sketch the derivation these results from a model field theory which has the same structure as the field theory that describes the quantum phase transition in disordered itinerant Heisenberg ferromagnets. In addition to demonstrating the origin of the breakdown of the loop expansion manifest in Eqs. (3), it will show that this breakdown is technically related to the field theory being non-renormalizable in a way that cannot be cured by introducing additional renormalization constants. This non-renormalizability is equivalent to the dangerous irrelevancy of w.

Let us consider a quantum field theory that couples diffusive excitations q to a ϕ^4-theory, with Gaussian propagators

$$\langle q_n(\boldsymbol{p})\, q_{-n}(-\boldsymbol{p})\rangle = \frac{G}{\boldsymbol{p}^2 + GH|\Omega_n|}\ , \tag{4a}$$

$$\langle \phi_n(\boldsymbol{p})\, \phi_{-n}(-\boldsymbol{p})\rangle = \frac{1}{t + a\boldsymbol{p}^2 + \frac{aC_1|\Omega_n|}{\boldsymbol{p}^2 + GH|\Omega_n|}}\ . \tag{4b}$$

The fields q and ϕ are functions of a wave vector $\boldsymbol{p}$ and a Matsubara frequency $\Omega_n = 2\pi T n$. The propagators in Eqs. (4a) and (4b) have structures appropriate for diffusive particle-hole excitations and paramagnons, respectively, in disordered metals. t is the distance from a critical point for the ϕ-excitation, and G, H, a, and C_1 are model parameters. Notice that the propagators contain the two time scales mentioned above. This will be crucial later. Motivated by the known properties of the ferromagnetic problem [11] we now assume that under renormalization, t, G, and C_1 are not singularly affected, while a and H acquire momentum and frequency dependences, respectively, that are determined by a set of coupled integral equations which read, at quantum criticality, $T = t = 0$, and with a momentum cutoff Λ,

$$a(\boldsymbol{k}) = a + \frac{C_2}{3}\int_0^\infty d\omega \int_{|\boldsymbol{k}|}^\Lambda dp\ \frac{p^{d-1}}{\left[p^2 + H(\omega)\omega\right]^3}\ . \tag{5a}$$

$$H(\Omega) = H + \frac{3C_2}{4}\int_0^\Lambda dp\ \frac{p^{d-1}}{a(\boldsymbol{p})p^4 + aC_1\Omega}\ . \tag{5b}$$

Here C_2 is another coupling constant.

While Eqs. (4,5) are sufficient to define our model, it is conceptually helpful to consider the structure of a field theory that generates these propagators. A Gaussian Landau-Ginzburg-Wilson (LGW) action consistent with Gaussian propagators given by Eqs. (4) has a structure

$$\mathcal{A}_0 = \int d\boldsymbol{x}\ \phi\,(t - a\,\partial_{\boldsymbol{x}}^2)\,\phi + \sqrt{T}\gamma_1 \int d\boldsymbol{x}\ \phi\, q + \int d\boldsymbol{x}\ q\,(-\partial_{\boldsymbol{x}}^2/G + H|\Omega_n|)\,q. \tag{6}$$

Here we consider the fields ϕ and q functions of position $\boldsymbol{x}$ and Ω_n, and we use a schematic notation that leaves out everything not needed for power counting, including sums over Matsubara frequencies and the distinction between powers of T and Ω_n. The coefficients t, a, G, and H are the same as in Eqs. (4), and $\gamma_1 = \sqrt{C_1 a/G}$. The renormalizations given by Eqs. (5) are produced by a non-Gaussian term with a structure

$$\mathcal{A}_{\text{int}} = \sqrt{T}\gamma_2 \int d\boldsymbol{x}\ \phi\, q^2, \tag{7}$$

with $\gamma_2 = \sqrt{GC_2/a}$. It gives rise to renormalizations of a and H by means of the diagrams shown in Fig. 1 [16].

The above model shares all qualitative features with the theory of disordered itinerant Heisenberg quantum ferromagnets that was considered in Refs. [11, 12]. In



FIG. 1: Diagrammatic contributions to $a(\boldsymbol{k})$ (a) and $H(\Omega)$ (b), respectively. The solid and dashed lines denote q and ϕ-propagators, respectively.

particular, the integral equations (5) are slightly simplified versions of those derived and solved in Refs. 12, 17. The prefactors in Eqs. (5) have been chosen such that the one-loop terms quantitatively agree with those obtained from the full integral equations.

Since the perturbation theory is given to all orders by the integral equations (5), one can determine the critical behavior by solving the latter. This was done in Ref. 12. In what follows we instead analyze the theory by standard RG techniques [6, 7, 8] order by order in a loop expansion. As we will see, this leads to important insights which one misses by solving the integral equations, and which transcend the specific ferromagnetic quantum phase transition problem that inspired our model.

We define the scale dimensions of a length L and frequency Ω or temperature T to be $[L] = -1$, $[\Omega] = [T] = z$, with z a so far undetermined dynamical critical exponent [18]. The scale dimensions of the fields ϕ and q we define such that a and G are dimensionless. We further define dimensionless coupling constants h, c_1, and c_2 by

$$H = \Lambda^{2-z} h \quad , \quad C_1 = \Lambda^{4-z} c_1 \quad , \quad C_2 = \Lambda^{2+\epsilon-z} c_2, \tag{8}$$

and derive RG flow equations for these quantities.

The properties of our model allow us to determine several flow equations and exponent relations exactly. Since G is not renormalized, the field q does not carry a field renormalization, which means that the diffusive propagators retains its Gaussian wave number dependence. ϕ, however, does carry a field renormalization, which gives rise to the critical exponent η defined by $a(\boldsymbol{p} \to 0) \propto |\boldsymbol{p}|^{-\eta}$. The fact that C_1 and t are not renormalized implies the exact flow equations

$$dc_1/d\ell = (4 - z - \eta) c_1 \quad , \quad dt/d\ell = (2 - \eta) t. \tag{9}$$

Both z and the correlation length exponent ν (defined by $t \propto b^{1/\nu}$) can therefore be expressed in terms of η and are given by Eq. (2b). The critical properties of the model are thus determined by only one independent exponent, e.g., η. Power counting shows that C_2 is not singularly renormalized either, so the flow equation for c_2 is also known exactly,

$$dc_2/d\ell = -(2 - \epsilon) c_2. \tag{10}$$

The flow equation for h, as well as η, need to be constructed in terms of a loop expansion.

Let us first consider the theory to zero-loop order. Since c_2 is irrelevant for all $d > 2$, Eq. (10), there is a Gaussian FP with $\eta = 0$, $z = 4$. This is the FP found by Hertz [2]. It is not stable, as was first shown in Ref. 4. In the present context, this can be seen as follows.

Consider the perturbation theory as generated by iterating the integral equations (5). For our present purposes it suffices to do the integrals in $d = 4$. With $U = C_2/aH$, the perturbation theory then has the structure

$$\begin{aligned} a(\boldsymbol{k})/a &= 1 + U F^{(1)}(\Lambda/|\boldsymbol{k}|) + U^2 \tilde{F}^{(2)}(\Lambda/|\boldsymbol{k}|) \\ &\quad + O(U^3), \end{aligned} \tag{11a}$$

$$\begin{aligned} H(\Omega)/H &= 1 + U\, G^{(1)}(\Lambda/\omega) + U^2\, G^{(2)}(\Lambda/\omega) \\ &\quad + O(U^3), \end{aligned} \tag{11b}$$

The proper expansion parameter is thus not C_2, but rather U, or its dimensionless counterpart $u = c_2/ah$, which for $d < 4$ is relevant with respect to the Gaussian FP with a scale dimension of ϵ, as can be seen from Eq. (8). The Gaussian FP is therefore stable only for $d > 4$, despite the irrelevancy of c_2, since h acts as a DIV [19].

To one-loop order, $F^{(1)}(x) = f_1 \ln x$ with $f_1 = 1/6$, and $G^{(1)}(x \to \infty) = g_1 \ln x + O(x^{-4})$ with $g_1 = 3/4$. Via standard methods [20] this yields Eqs. (1,2).

At two-loop order, $\tilde{F}^{(2)}(x \to \infty) = \tilde{f}_{2,2} \ln^2 x + O(\ln x)$ with $\tilde{f}_{2,2} = -3/32$, and $G^{(2)}(x \to \infty) = g_{2,2} \ln^2 x + g_{2,1} \ln x + O(1)$ with $g_{2,2} = -1/16$, $g_{2,1} = 1/32$. It is obvious, in either the momentum-shell or the field-theoretic RG methods, that the coefficients of the two-loop $\ln^2$-terms must not be independent of the coefficients of the one-loop ln-terms if the theory is to be renormalizable (in the latter method), or lead to scale-independent flow equations and critical exponents (in either method). A straightforward analysis shows that the condition is $\tilde{f}_{2,2} = g_{2,2} = -f_1 g_1/2$. This condition is obeyed by $g_{2,2}$, but not by $\tilde{f}_{2,2}$. This problem is germane to the structure of the perturbation theory and cannot be cured by introducing additional renormalization constants (within the field-theoretic method). To see its origin, let us analyze the contributions to the $\ln^2$-terms.

Equation (5b) for $H(\Omega)$ in our model has the same structure as insertions in a classical theory, since there is no frequency integral. The one-loop correction to $a(\boldsymbol{p})$ is linear in $\ln(\Lambda/p) = \ln \Lambda - \ln p$ with $p = |\boldsymbol{p}|$. Both of these terms contribute to the $\ln^2 \Lambda$ contribution to $H(\Omega)$, with relative prefactors of 1 and $-1/2$, respectively. This structure guarantees that $g_{2,2} = -f_1 g_1/2$. The one-loop contribution to $H(\Omega)$, on the other hand, is linear in $\ln(\Lambda/\Omega^{1/4})$, which leads to contributions to the $\ln^2 \Lambda$ term in $a(\boldsymbol{k})$ with relative prefactors of 1 and $-1/4$, respectively. As a result, one has $\tilde{f}_{2,2} = -(1-1/4) f_1 g_1 = -3 f_1 g_1/4$. This is a direct consequence of the existence of two time scales: If one replaced the Ω in the integrand in Eq. (5b) by Ω^2, which would result in a model with only one time scale, $\Omega \sim \boldsymbol{p}^2$, one would get $\tilde{f}_{2,2} = g_{2,2} = -f_1 g_1/2$.

These observations suggest the following interpretation of this failure of the renormalization procedure. After scaling the integration variable ω in Eq. (5a) with p^2/H, the one-loop correction to $H(\omega)$ is proportional to $\ln(\Lambda^4 H/C_1\omega p^2)$. Writing this as $\ln(\Lambda^4/p^4) - \ln\omega + \ln(Hp^2/C_1)$, we can write $\tilde{F}^{(2)}$ in Eq. (11a) as

$$\tilde{F}^{(2)}(\Lambda/|\boldsymbol{k}|) = F^{(2)}(\Lambda/|\boldsymbol{k}|) - \frac{1}{32}\int_{|\boldsymbol{k}|}^{\Lambda} dp\,\frac{1}{p}\,\ln w(p), \quad (12)$$

where $F^{(2)}(x \to \infty) = -(\ln x)^2/16 + O(\ln x)$, and $w(p) = Hp^2/C_1 = (h/c_1)(p/\Lambda)^2$. This leads to the two-loop flow equations (3), where $w = w(p = \Lambda) = h/c_1$. The FP value of the irrelevant variable w is $w^* = 0$. Inserted into the u-flow equation, this yields an infinite coefficient at two-loop order. We also see that the physical origin of this problem is the second (diffusive) time scale, which is reflected in the scale dimension of $w \propto h$.

The following picture now emerges. h is a DIV since the loop expansion is in powers of $u = c_2/ah$. At one-loop order, this makes the Gaussian FP unstable, but one has the seemingly sensible FP discussed in connection with Eqs. (1,2). The two-loop term, however, goes as $h^{-2}\ln h$ rather than just as $1/h^2$, and this leads to a breakdown of the loop expansion. If one eliminates w by solving Eq. (3b) and substituting the result in Eq. (3a), one obtains scale dependent flow equations and critical exponents. As we have seen, this is a consequence of the presence of two time scales in conjunction with the structure of the perturbation theory for our model. It is therefore not possible to analyze this field theory by means of standard RG methods. At higher order in the loop expansion, higher powers of $\ln w$ will appear. Clearly, a resummation to all orders is needed in order to determine the critical behavior. Indeed, a solution of the integral equations, which achieve such a resummation, shows that the asymptotic critical behavior consists of power laws multiplied by log-normal terms [12].

In summary, we have used the quantum ferromagnetic transition in disordered metals to show that one cannot necessarily trust low-order perturbative RG results, even in the absence of any indications that the perturbative FP one has found might not be stable. The mechanism we have discussed involves a DIV and is rather general, and is thus expected to be operative at other phase transitions as well. For instance, our analysis casts doubt on a recent one-loop calculation for the clean quantum ferromagnetic problem [21]. We note, however, that the presence of multiple time scales, while necessary, is not sufficient to produce the effect. For instance, at the liquid-gas critical point mentioned in the introduction, the analog of the irrelevant variable w is not dangerous [22].

This work was supported by the NSF under grant Nos. DMR-01-32555 and DMR-01-32726, and by a fellowship of the Deutsche Forschungsgemeinschaft (J.R.). Part of this work was performed at the Aspen Center for Physics.

---

[1] See, e.g., P. C. Hohenberg and B. I. Halperin, Rev. Mod. Phys. **49**, 435 (1977).

[2] J. Hertz, Phys. Rev. B **14**, 1165 (1976).

[3] A. J. Millis, Phys. Rev. B **48**, 7183 (1993).

[4] T. R. Kirkpatrick and D. Belitz, Phys. Rev. B **53**, 14364 (1996).

[5] For a recent review, see, D. Belitz, T. R. Kirkpatrick, and T. Vojta, cond-mat/0403182.

[6] K. G. Wilson and J. Kogut, Phys. Rep. **12**, 75 (1974).

[7] S.-K. Ma, *Modern Theory of Critical Phenomena* (Benjamin, Reading, MA, 1976).

[8] See, e.g., J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena* (Oxford University Press, Oxford, 1996).

[9] S. Weinberg, *The Quantum Theory of Fields*, vol. II (Cambridge University, Cambridge, 1996).

[10] M. E. Fisher, in *Advanced Course on Critical Phenomena*, edited by F. W. Hahne (Springer, Berlin, 1983), p. 1.

[11] D. Belitz, T. R. Kirkpatrick, M. T. Mercaldo, and S. L. Sessions, Phys. Rev. B **63**, 174427 (2001).

[12] D. Belitz, T. R. Kirkpatrick, M. T. Mercaldo, and S. L. Sessions, Phys. Rev. B **63**, 174428 (2001).

[13] An analogous irrelevant variable appears at the liquid-gas critical point, but it is not dangerous [22].

[14] For a review, see, J. R. Dorfman, T. R. Kirkpatrick, and J. V. Sengers, Ann. Rev. Phys. Chem. A **45**, 213 (1994).

[15] According to a tree-level analysis, $z = 4$ implies an upper critical dimension $d_c^+ = 0$ [2]. This is fallacious, however, as can be seen from the fact that y becomes relevant with respect to the mean-field fixed point for $d < 4$. This is the first manifestation of the dangerous irrelevancy of w, as we will see below.

[16] Several features of the integral equations (5) are not obvious from the schematic action, Eqs. (6,7), e.g., the fact that the correction to H contains no frequency integral. In order to see this, and other points, one needs to acknowledge that the diffusive particle-hole excitations q actually depend on two fermionic frequencies rather than one bosonic one, and one also needs to consider a q^4-vertex [11, 12]. Here we suppress these technicalities, which only distract from our main message, consider Eqs. (4,5) the definition of our model, and use the schematic action only to explain the structure of the RG analysis.

[17] T. R. Kirkpatrick and D. Belitz, Phys. Rev. B **45**, 3187 (1992).

[18] We choose to introduce only one dynamical critical exponent z, despite the fact that there are two time scales in the problem. The second one is implicit in the scale dimension of H. This is same procedure as in Ref. 22.

[19] We use the term "dangerous irrelevant variable" in a very general sense: The loop expansion parameter itself depends singularly on the irrelevant variable h, not just some scaling function.

[20] To one-loop order one can use either a momentum-shell RG, or a field-theoretic one, to derive these results. At two-loop order, only the field-theoretic method is practical, as the shell integrals become too complicated.

[21] T. R. Kirkpatrick and D. Belitz, Phys. Rev. B **67**, 024419 (2003).

[22] C. DeDominicis and L. Peliti, Phys. Rev. B **18**, 353 (1978).